

## **IMPACT OF QUARRY DUST POLLUTION ON FOLIAR EPIDERMIS OF FIVE SPECIES GROWING NEAR STONE CRUSHING INDUSTRY**

**D. PARAMESHA NAIK, USHAMALINI AND R.K. SOMASHEKAR**

Department of Environmental Sciences, Bangalore University,  
Bangalore 560 056, Karnataka, India

**Key words :** Quarry dust, Anatomical variation, Air pollution .

### **ABSTRACT**

---

---

**Quarry dust is known to alter the stomatal index of most of the plants studies. But in some plants the stomatal index remains unaffected or undergo little changes. Study carried out on selected plant species growing in the vicinity revealed foliar epidermal responses to quarry dust pollution which differed from species to species. The data obtained from the seasonal study revealed the impact of crushing activity, which has resulted in the alternation of anatomical features in various plant species.**

---

---

### **INTRODUCTION**

Due to unscientific and unplanned crushing activity large scale fine dust particles are released in to the atmosphere which has the potential to alter the anatomical characteristics of the plants species growing in the near vicinity. The dust particles fall heavily on the near by vegetation which bring about changes in the anatomical features. Foliar epidermal responses to suspended particulate matter are worked out by various scientists (Gouse *et al.* 1972, Sharma *et al.* 1973, 1975, Gouse, *et al.* 1978, Rajachidambaram *et al.* ). Recent attempts have shown that epidermis responds to various pollutants and therefore can be employed as a biological indicator to measure the levels of atmospheric pollution (Sharma, *et al.* 1973). The present work deals with the foliar epidermal responses of different species growing near stone crushing industry.

**Table 1.** Foliar epidermal variations of five different plant species from polluted and unpolluted quarry area.

Slno	Plant species	(Values are Mean $\pm$ S.D.)											
		Calotropis gigantea		Eucalyptus globulus		Psidium guajava		Muntingia calabura		Croton sparsiflorus			
		UE	LE	UE	LE	UE	LE	UE	LE	UE	LE		
Stomal index	P	4.6 $\pm$ 1.12	6.8 $\pm$ 2.12	7.1 $\pm$ 1.17	9.8 $\pm$ 1.74	8.0 $\pm$ 2.16	9.2 $\pm$ 2.16	7.9 $\pm$ 2.10	9.1 $\pm$ 1.73	6.8 $\pm$ 2.1	8.7 $\pm$ 1.35		
1 MM <sup>2</sup>	U	8.4 $\pm$ 2.16	7.2 $\pm$ 10.42	6.7 $\pm$ 1.72	7.9 $\pm$ 1.67	8.2 $\pm$ 1.36	9.5 $\pm$ 1.92	7.5 $\pm$ 1.32	8.9 $\pm$ 2.16	6.9 $\pm$ 1.1	9.4 $\pm$ 12.0		
Length of guard cells in $\mu$	P	215.5 $\pm$ 24.7	215.0 $\pm$ 22.6	228.1 $\pm$ 11.2	229.0 $\pm$ 10.9	123.2 $\pm$ 12.3	123.0 $\pm$ 12.10	141.0 $\pm$ 40.2	140.0 $\pm$ 40.2	210.2 $\pm$ 18.2	211.0 $\pm$ 18.2		
	U	256.2 $\pm$ 15.1	256.0 $\pm$ 25.5	220.0 $\pm$ 20.7	223.2 $\pm$ 20.0	120.5 $\pm$ 11.12	12.0 $\pm$ 11.3	139. $\pm$ 21.2	139.0 $\pm$ 21.2	221.1 $\pm$ 22.6	220.0 $\pm$ 12.5		
Width of guard cell in $\mu$	P	52.5 $\pm$ 21.5	65.6 $\pm$ 14.9	63.8 $\pm$ 7.0	63.8 $\pm$ 7.0	60.8 $\pm$ 5.72	60.8 $\pm$ 5.72	52.12 $\pm$ 5.12	52.10 $\pm$ 5.11	60.4 $\pm$ 10.5	63.0 $\pm$ 13.02		
	U	80.5 $\pm$ 13.8	82.5 $\pm$ 13.1	60.7 $\pm$ 6.11	60.7 $\pm$ 6.10	61.4 $\pm$ 6.27	61.4 $\pm$ 6.27	49.18 $\pm$ 5.10	49.18 $\pm$ 5.10	62.5 $\pm$ 8.16	62.5 $\pm$ 8.19		
Length stomata pore in $\mu$	P	141.5 $\pm$ 30.6	141.6 $\pm$ 18.1	154.6 $\pm$ 20.2	154.6 $\pm$ 20.2	98.0 $\pm$ 7.60	101.2 $\pm$ 9.52	84.6 $\pm$ 10.2	84.0 $\pm$ 10.0	115.9 $\pm$ 10.2	14.9 $\pm$ 10.18		
	U	155.4 $\pm$ 20.7	156.0 $\pm$ 20.7	150.2 $\pm$ 5.28	150.2 $\pm$ 5.27	95.9 $\pm$ 5.70	95.9 $\pm$ 6.23	80.7 $\pm$ 11.3	80.5 $\pm$ 11.0	122.6 $\pm$ 18.8	122.2 $\pm$ 18.0		
Width of stomatal pore in $\mu$	P	19.7 $\pm$ 5.72	19.7 $\pm$ 5.36	64.0 $\pm$ 6.80	63.0 $\pm$ 8.22	23.2 $\pm$ 6.21	23.0 $\pm$ 6.23	20.8 $\pm$ 6.28	20.7 $\pm$ 6.28	24.8 $\pm$ 8.70	23.7 $\pm$ 6.70		
	U	26.5 $\pm$ 5.2	27.7 $\pm$ 6.4	60.7 $\pm$ 8.42	60.7 $\pm$ 8.39	24.8 $\pm$ 4.94	24.6 $\pm$ 4.90	20.4 $\pm$ 6.46	20.3 $\pm$ 6.40	20.1 $\pm$ 5.28	20.1 $\pm$ 5.28		

Note: P - Polluted, UP - Unpolluted, UE - Upper epidermis, LE - Lower epidermis

## MATERIALS AND METHODS

Leaves of *Calotropis gigantea*, *Eucalyptus guajava*, *Muntingia calabura* and *Croton sparsiflorus* were collected near stone crushing site. 25 samples were collected randomly from each site. For herbs 5th leaf from base of the plants of uniform size were collected. For herbs, shrubs or trees leaves from 2 mts height were collected. Leaves were immediately fixed in the F.A.A. solution. For unpolluted controls leaves of same species were collected 25km away from the quarry area. Stomatal index was calculated as per Salisbury, (1928). Epidermal peels of the leaves were obtained by hot sulphuric acid treatment (Gouse *et al.* 1972).

## RESULTS AND DISCUSSION

Stomatal index, length and width of guard cells, length and width of stomatal pore were decreased in *Calotropis gigantea* species in both upper and lower epidermis. In *Eucalyptus globules*, *Psidium guajava* and *Muntingia calabura* plant species, stomatal index was more in polluted leaves, but the other parameters showed a decrease. In *Croton sparsiflorus* species polluted and inpolluted leaf epidermis showed littel variations. Similar results were reported by Krishnamurthy *et al.* (1980). The results obtained depicts the gloomy picture of the crushing activity and its impact on stomatal index and other epidermal variations of the plant species studied.

## REFERENCES

- Ghose, A.K.M. and Khan, A.U. 1978. *Proc. Int. Sym. Env. Hyderabad.* 41- 43.  
 Ghose, A.K.M. and Yunus, M. 1972. *Stain Technol.* 47 : 322 - 324.  
 Krishnamurthy, K.V. and Raja Chidambaram, 1980. *C. Biology.* 2 : 8 -13.  
 Salisbury, E.J.-Phil, 1928. *Trans Royal. Soc. (London)* 216 : 1- 65.  
 Sharma, G.K. and Butler, 1973. *Jr. of Environ. Poll.* 5 : 267 - 293.  
 Sharma, G.K. and Butler, 1975. *J. Ann. Bot.* 39 : 1087 - 1090.

## ENCYCLOPEDIA OF ENVIRONMENTAL POLLUTION AND CONTROL

R.K.Trivedy

**(Vol. 1)** • Acid rain • Air pollution: sources, abundance and new challenges • Adsorbents for the treatment of wastewaters • Air pollution and plants • Air pollution and control • Air pollution control for thermal power stations • Algae in water supply systems and their control • Anaerobic processes • Bacterial pollution and water disinfection procedures • Biological problems in industrial cooling and control • Biomonitoring of water pollution • Conventional methods of waste water treatment • Designing wastewater treatment plants • Effect of air pollution on Man and Property • Environmental software • Gaseous pollution control • Green belt for pollution abatement • Eutrophication • Hazardous waste management • Heavy metal pollution: Origin, impact and remedies.

**(Vol. 2)** • Incineration for waste management • Indian legislation for pollution control • Indoor air pollution • Industry and water pollution • Land treatment of waste water • Marine pollution • Nitrogen and phosphorus removal from wastewater • Noise pollution and control • Oil pollution-treatment of oil wastes and oil spillages • Ozone layer protection • Pesticide transformation and metabolism • Physico-chemical methods in monitoring water pollution • Pollution control by membrane technology • Pollution from thermal power plants • Risk assessment and disaster management in chemical industries • River ecology and water pollution • Simple and rapid enzymatic methods for environmental monitoring of some heavy metals in water and agricultural products • Solid waste utilization • Siting of industries- An environmental outlook • Sources of water pollution and water quality parameters • Vehicular pollution in India • Treatment and disposal of domestic sewage • Waste recycling -The key to cleaner technologies • Waste water treatment theory and practice of biological sludge production • Water pollution and Fisheries.

**PAGES 900, Price Rs. 1500.00 US\$ 100.00, Paperback Rs.700.00  
US\$ 50.00**

**All Enquiries, Orders & Correspondence to  
ENVIRO MEDIA - 2nd Floor, Rohan Heights,  
Karad 415 110 (India)  
Tel : 912164 -220369, 020 - 27210103,  
Email - rktem@pn3.vsnl.net.in**